When the property of having a π-tree is preserved by products*

Mikhail Patrakeev †

arXiv:1611.08870v1 [math.GN] 27 Nov 2016

Abstract

We find sufficient conditions under which the product of spaces that have a π-tree also has a π-tree. These conditions give new examples of spaces with a π-tree: every at most countable power of the Sorgenfrey line and every at most countable power of the irrational Sorgenfrey line has a π-tree. Also we show that if a space has a π-tree, then its product with the Baire space, with the Sorgenfrey line, and with the countable power of the Sorgenfrey line also has a π-tree.

1 Introduction

We study topological spaces that have a π-tree; the notion of a π-tree was introduced in [1] and is equivalent [1, Remark 11] to the notion of a Lusin π-base, which was introduced in [2]. The Sorgenfrey line $\mathcal{R}_\mathcal{S}$ and the Baire space $\mathcal{N}$ (that is, ${}^\omega\omega$ with the product topology) are examples of spaces with a π-tree [2]. Every space that has a π-tree shares many good properties with the Baire space. One reason for this is expressed in Lemmas 6 and 9, another two are the following: If a space X has a π-tree, then X can be mapped onto $\mathcal{N}$ by a continuous one-to-one map [2] and also X can be mapped onto $\mathcal{N}$ by a continuous open map [2] (hence X can be mapped by a continuous open map onto an arbitrary Polish space, see [3] or [4, Exercise 7.14]). Every space that has a π-tree also has a countable π-base, see Lemma 7.

In this paper we study the following question: When does the product of spaces that have a π-tree also have a π-tree? We find several kinds of conditions (see Theorems 14, 17 and Corollary 15) under which an at most countable product of spaces that have a π-tree also has a π-tree. We consider only at most countable products because an uncountable product of spaces that have a π-tree has an uncountable pseudocharacter, therefore it has no π-tree (see [5, statement 5.3.b] and Lemmas 6, 7, 9).

The above results give new examples of spaces that have a π-tree, see Section 7. For instance, Corollary 25 assirts that if $1 \leqslant |A| \leqslant \omega$ and for each $\alpha \in A$,

$$\text{either} \qquad X_\alpha = \mathcal{N} \qquad \text{or} \qquad X_\alpha \subseteq \mathcal{R}_\mathcal{S} \ \text{ with } \ \mathcal{R}_\mathcal{S} \setminus X_\alpha \ \text{ at most countable,}$$

then the product $\prod_{\alpha\in A} X_\alpha$ has a π-tree. In particular, the powers ${\mathcal{R}_\mathcal{S}}^n$ and ${\mathcal{I}_\mathcal{S}}^n$ ($\mathcal{I}_\mathcal{S}$ denotes the irrational Sorgenfrey line $\mathcal{R}_\mathcal{S} \setminus \mathbb{Q}$) have a π-tree for all natural $n \geqslant 1$, and the powers ${\mathcal{R}_\mathcal{S}}^\omega$ and ${\mathcal{I}_\mathcal{S}}^\omega$ also have a π-tree. (Note that no finite power of the irrational Sorgenfrey line is homeomorphic to finite power of the Sorgenfrey line [6].) Other examples of spaces with a π-tree can be obtained by using Corollary 27, which says that if a space X has a π-tree, then the products $X \times \mathcal{N}$, $X \times \mathcal{R}_\mathcal{S}$, and $X \times {\mathcal{R}_\mathcal{S}}^\omega$ also have a π-tree.

*2010 Mathematics Subject Classification: Primary 54E99; Secondary 54H05. Keywords: the Sorgenfrey line, the Baire space, Souslin scheme, Lusin scheme, open sieve, Lusin pi-base, pi-tree, foliage tree, the foliage hybrid operation, product of topological spaces.

†Krasovskii Institute of Mathematics and Mechanics of UB RAS, 620990, 16 Sofia Kovalevskaya street, Yekaterinburg, Russia and Ural Federal University, 620002, 19 Mira street, Yekaterinburg, Russia; patrakeev@mail.ru

2 Notation and terminology

We use standard set-theoretic notation from [7, 8]. In particular, each ordinal is equal to the set of smaller ordinals, ω = the set of natural numbers = the set of finite ordinals = the first limit ordinal = the first infinite cardinal, and $n = \{0, \dots, n-1\}$ for all $n \in \omega$. A *space* is a topological space; we use terminology from [9] when we work with spaces. Also we use the following notations:

Terminology 1. The symbol $:=$ means "equals by definition"; the symbol $:\longleftrightarrow$ is used to show that an expression on the left side is an abbreviation for expression on the right side;

- $x \subset y \quad :\longleftrightarrow \quad x \subseteq y$ and $x \neq y$;
- $A \equiv \bigsqcup_{\lambda\in\Lambda} B_\lambda \quad :\longleftrightarrow \quad A = \bigcup_{\lambda\in\Lambda} B_\lambda$ and $\forall \lambda, \lambda' \in \Lambda\, [\lambda \neq \lambda' \to B_\lambda \cap B_{\lambda'} = \varnothing]$;
- $[A]^\kappa := \big\{B \subseteq A : |B| = \kappa\big\}, \quad [A]^{<\kappa} := \big\{B \subseteq A : |B| < \kappa\big\}$ (here κ is a cardinal);
- γ has the FIP $\quad :\longleftrightarrow \quad \forall \delta \in [\gamma]^{<\omega} \setminus \{\varnothing\} \left[\bigcap \delta \neq \varnothing\right]$ (FIP means finite intersection property);
- $\operatorname{cofin} A := \big\{A \setminus F : F \in [A]^{<\omega}\big\}$;
- $\operatorname{nbhds}(p, X) :=$ the set of (not necessarily open) neighbourhoods of point p in space X;
- $f \restriction A :=$ the restriction of function f to A;
- $\gamma \gg \delta \quad :\longleftrightarrow \quad \gamma$ *π-refines* $\delta \quad :\longleftrightarrow \quad \forall D \in \delta \setminus \{\varnothing\}\ \exists G \in \gamma \setminus \{\varnothing\}\ [G \subseteq D]$.

When we work with (transfinite) sequences, we use the following notations:

Terminology 2. Suppose $n \in \omega$ and s, t are sequences; that is, s and t are functions whose domain is an ordinal.

- $\operatorname{length} s :=$ the domain of s;
- note that $s \subseteq t$ iff $\operatorname{length} s \leqslant \operatorname{length} t$ and $s = t \restriction \operatorname{length} s$;
- $\langle r_0, \dots, r_{n-1}\rangle :=$ the sequence r such that $\operatorname{length} r = n$ and $r(i) = r_i$ for all $i \in n$;
- $\langle\rangle :=$ the sequence of length 0;
- $\langle r_0, \dots, r_{n-1}\rangle^\frown\langle s_0, \dots, s_{m-1}\rangle := \langle r_0, \dots, r_{n-1}, s_0, \dots, s_{m-1}\rangle$;
- ${}^BA :=$ the set of functions from B to A; in particular, ${}^0A = \big\{\langle\rangle\big\}$;
- ${}^{<\alpha}A := \bigcup_{\beta\in\alpha} {}^\beta A$ (here α is an ordinal).

Also we work with partial orders and then we use the following terminology:

Terminology 3. Suppose $\mathcal{P} = (Q, \lhd)$ is a strict partial order; that is, $\lhd$ is irreflexive and transitive on Q. Let $x, y \in Q$ and $A \subseteq Q$.

- $\operatorname{nodes}\mathcal{P} = \operatorname{nodes}(Q, \lhd) := Q$;
- $x <_\mathcal{P} y \quad :\longleftrightarrow \quad x \lhd y$;
- $x \leqslant_\mathcal{P} y \quad :\longleftrightarrow \quad x <_\mathcal{P} y$ or $x = y$;
- $x\overset{\circ}{\uparrow}_\mathcal{P} := \{v \in \operatorname{nodes}\mathcal{P} : v <_\mathcal{P} x\}, \quad x\overset{\circ}{\downarrow}_\mathcal{P} := \{v \in \operatorname{nodes}\mathcal{P} : v >_\mathcal{P} x\}$;
- $x\overset{\bullet}{\uparrow}_\mathcal{P} := \{v \in \operatorname{nodes}\mathcal{P} : v \leqslant_\mathcal{P} x\}, \quad x\overset{\bullet}{\downarrow}_\mathcal{P} := \{v \in \operatorname{nodes}\mathcal{P} : v \geqslant_\mathcal{P} x\}$;
- $A\top_\mathcal{P} := \bigcup\{v\overset{\bullet}{\uparrow}_\mathcal{P} : v \in A\}, \quad A\bot_\mathcal{P} := \bigcup\{v\overset{\bullet}{\downarrow}_\mathcal{P} : v \in A\}$;
- $\operatorname{sons}_\mathcal{P}(x) := \{s \in \operatorname{nodes}\mathcal{P} : x <_\mathcal{P} s$ and $x\overset{\circ}{\downarrow}_\mathcal{P} \cap s\overset{\circ}{\uparrow}_\mathcal{P} = \varnothing\}$;
- A is a *chain* in $\mathcal{P} \quad :\longleftrightarrow \quad \forall v, w \in A\, [v \leqslant_\mathcal{P} w$ or $v >_\mathcal{P} w]$;
- $\mathcal{P}$ has *bounded chains* $\quad :\longleftrightarrow$
  for each nonempty chain C in $\mathcal{P}$ there is $v \in \operatorname{nodes}\mathcal{P}$ such that $C \subseteq v\overset{\bullet}{\uparrow}_\mathcal{P}$;
- $\max\mathcal{P} := \{m \in \operatorname{nodes}\mathcal{P} : m\overset{\circ}{\downarrow}_\mathcal{P} = \varnothing\}, \quad \min\mathcal{P} := \{m \in \operatorname{nodes}\mathcal{P} : m\overset{\circ}{\uparrow}_\mathcal{P} = \varnothing\}$;
- $0_\mathcal{P} :=$ the node such that $(0_\mathcal{P})\overset{\bullet}{\downarrow}_\mathcal{P} = \operatorname{nodes}\mathcal{P}$ (here $\mathcal{P}$ is a partial order that has such node).

When a partial order is a (set-theoretic) tree, we use the following terminology:

Terminology 4. Suppose $\mathcal{T}$ is a tree; that is, $\mathcal{T}$ is a strict partial order such that for each $x \in \mathsf{nodes}\,\mathcal{T}$, the set $x{\restriction_{\mathcal{T}}}$ is well-ordered by $<_{\mathcal{T}}$. Let $x \in \mathsf{nodes}\,\mathcal{T}$, let α be an ordinal, and let κ be a cardinal.

- $\mathsf{height}_{\mathcal{T}}(x) :=$ the ordinal isomorphic to $(x{\restriction_{\mathcal{T}}}, <_{\mathcal{T}})$;
- $\mathsf{level}_{\mathcal{T}}(\alpha) := \big\{v \in \mathsf{nodes}\,\mathcal{T} : \mathsf{height}_{\mathcal{T}}(v) = \alpha\big\}$;
- $\mathsf{height}\,\mathcal{T} :=$ the minimal ordinal β such that $\mathsf{level}_{\mathcal{T}}(\beta) = \varnothing$;
- B is a *branch* in $\mathcal{T}$ $:\longleftrightarrow$ B is a $\subseteq$-maximal chain in $\mathcal{T}$;
- $\mathcal{T}$ is *κ-branching* $:\longleftrightarrow$ $\forall v \in \mathsf{nodes}\,\mathcal{T} \setminus \mathsf{max}\,\mathcal{T}\ \big[|\mathsf{sons}_{\mathcal{T}}(v)| = \kappa\big]$.

Finally, we work with foliage trees, which where introduced in [1]. Recall that a *foliage tree* is a pair $\mathbf{F} = (\mathcal{T}, l)$ such that $\mathcal{T}$ is a tree and l is a function with $\mathsf{domain}\,l = \mathsf{nodes}\,\mathcal{T}$. For each $x \in \mathsf{nodes}\,\mathcal{T}$, the $l(x)$ is called the *leaf* of $\mathbf{F}$ at node X and is denoted by $\mathbf{F}_x$; the tree $\mathcal{T}$ is called the *skeleton* of $\mathbf{F}$ and is denoted by $\mathsf{skeleton}\,\mathbf{F}$. We adopt the following convention: If $\mathbf{F}$ is a foliage tree and $\bullet$ is a notation that can be applied to a tree, then $\bullet(\mathbf{F})$ is an abbreviation for $\bullet(\mathsf{skeleton}\,\mathbf{F})$; for example, $x <_{\mathbf{F}} y$ stands for $x <_{\mathsf{skeleton}\,\mathbf{F}} y$. Also we use the following terminology:

Terminology 5. Suppose $\mathbf{F}$ is a foliage tree, $v \in \mathsf{nodes}\,\mathbf{F}$, $A \subseteq \mathsf{nodes}\,\mathbf{F}$, X is a space, α is an ordinal, and κ is a cardinal.

- $\mathsf{flesh}\,\mathbf{F} := \bigcup\{\mathbf{F}_x : x \in \mathsf{nodes}\,\mathbf{F}\}$;
- $\mathsf{flesh}_{\mathbf{F}}(A) := \bigcup\{\mathbf{F}_x : x \in A\}$;
- $\mathsf{shoot}_{\mathbf{F}}(v) := \big\{\mathsf{flesh}_{\mathbf{F}}(C) : C \text{ is a cofinite subset of } \mathsf{sons}_{\mathbf{F}}(v)\big\}$;
- $\mathsf{scope}_{\mathbf{F}}(a) := \{x \in \mathsf{nodes}\,\mathbf{F} : \mathbf{F}_x \ni a\}$;
- $\mathbf{F}$ has *nonempty leaves* $:\longleftrightarrow$ $\forall x \in \mathsf{nodes}\,\mathbf{F}\ [\mathbf{F}_x \neq \varnothing]$;
- $\mathbf{F}$ is *nonincreasing* $:\longleftrightarrow$ $\forall x, y \in \mathsf{nodes}\,\mathbf{F}\ [y \geqslant_{\mathbf{F}} x \to \mathbf{F}_y \subseteq \mathbf{F}_x]$;
- $\mathbf{F}$ has *strict branches* $:\longleftrightarrow$
  $\mathsf{nodes}\,\mathbf{F} \neq \varnothing$ and for each branch B in $\mathbf{F}$, the $\bigcap_{x\in B}\mathbf{F}_x$ is a singleton;
- $\mathbf{F}$ is *locally strict* $:\longleftrightarrow$ $\forall x \in \mathsf{nodes}\,\mathbf{F} \setminus \mathsf{max}\,\mathbf{F}\ [\mathbf{F}_x \equiv \bigsqcup_{s\in\mathsf{sons}_{\mathbf{F}}(x)}\mathbf{F}_s]$;
- $\mathbf{F}$ is *open* in X $:\longleftrightarrow$ $\forall z \in \mathsf{nodes}\,\mathbf{F}\ [\mathbf{F}_z$ is an open subset of $X]$;
- $\mathbf{F}$ is a *foliage α,κ-tree* $:\longleftrightarrow$ $\mathsf{skeleton}\,\mathbf{F}$ is isomorphic to the tree $({}^{<\alpha}\kappa, \subset)$;
- $\mathbf{F}$ is a *Baire foliage tree* on X $:\longleftrightarrow$ $\mathbf{F}$ is an open in X locally strict foliage ω,ω-tree with strict branches and such that $\mathbf{F}_{0_{\mathbf{F}}} = X$;
- $\mathbf{F}$ *grows into* X $:\longleftrightarrow$ $\forall p{\in}X\ \forall U{\in}\mathsf{nbhds}(p,X)\ \exists z{\in}\mathsf{scope}_{\mathbf{F}}(p)\ \big[\mathsf{shoot}_{\mathbf{F}}(z) \gg \{U\}\big]$;
- $\mathbf{F}$ is a *π-tree* on X $:\longleftrightarrow$ $\mathbf{F}$ is a Baire foliage tree on X and $\mathbf{F}$ grows into X;
- $\mathbf{S} :=$ the *standard foliage tree* of ${}^{\omega}\omega :=$ the foliage tree such that
  - $\mathsf{skeleton}\,\mathbf{S} := ({}^{<\omega}\omega, \subset)$ and
  - $\mathbf{S}_x := \{p \in {}^{\omega}\omega : x \subseteq p\}$ for every $x \in {}^{<\omega}\omega$;
- $\mathcal{N} :=$ the *Baire space* $:=$ the space $({}^{\omega}\omega, \tau_{\mathcal{N}})$, where $\tau_{\mathcal{N}}$ is the Tychonoff product topology with ω carrying the discrete topology.

Lemma 6 (Lemma 13 in [1])**.**

(a) $\{\mathbf{S}_x : x \in {}^{<\omega}\omega\}$ *is a base for* $\mathcal{N}$.

(b) $\mathbf{S}$ *is a π-tree on* $\mathcal{N}$.

(c) $\mathbf{S}$ *is a Baire foliage tree on a space* $({}^{\omega}\omega, \tau)$ iff $\tau \supseteq \tau_{\mathcal{N}}$.

Lemma 7. *If* $\mathbf{F}$ *is a π-tree on a space* X*, then*

➢ $\{\mathbf{F}_v : v \in \mathsf{nodes}\,\mathbf{F}\}$ *is a countable* π*-base for* X,
➢ *each* $\mathbf{F}_v$ *is closed-and-open in* X, *and*
➢ $\bigcap\{\mathbf{F}_v : \mathbf{F}_v \ni p\} = \{p\}$ *for all* $p \in X$. □

3 New notions: isomorphism and spectrum

The notion of isomorphism between foliage trees allows to simplify proofs (see the proof of Theorem 14) in the following way: When we have a π-tree $\mathbf{F}$ on a space X, we may (by using (c) of Lemma 9 and (c) of Lemma 6) assume "without loss of generality" that $\mathbf{F} = \mathbf{S}$ and $X = ({}^{\omega}\omega, \tau)$ with $\tau \supseteq \tau_{\mathcal{N}}$.

Definition 8. An **isomorphism** between foliage trees $\mathbf{F}$ and $\mathbf{G}$ is a pair (φ, ψ) such that

➢ φ is an order isomorphism from $\mathsf{skeleton}\,\mathbf{F}$ onto $\mathsf{skeleton}\,\mathbf{G}$,
➢ ψ is a bijection from $\mathsf{flesh}\,\mathbf{F}$ onto $\mathsf{flesh}\,\mathbf{G}$, and
➢ $\psi[\mathbf{F}_x] = \mathbf{G}_{\varphi(x)}$ for all $x \in \mathsf{nodes}\,\mathbf{F}$.

Lemma 9. *Suppose that* $\mathbf{F}$ *is a foliage tree and* X *is a space.*

(a) $\mathbf{F}$ *is a locally strict foliage* ω,ω*-tree with strict branches* iff
$\mathbf{F}$ *is isomorphic to* $\mathbf{S}$.

(b) $\mathbf{F}$ *is a Baire foliage tree on* X iff
there exist an isomorphism (φ, ψ) *between* $\mathbf{F}$ *and* $\mathbf{S}$ *and a topology* τ *on* ${}^{\omega}\omega$ *such that*
➢ ψ *is a homeomorphism from* X *onto* $({}^{\omega}\omega, \tau)$ *and*
➢ $\mathbf{S}$ *is a Baire foliage tree on* $({}^{\omega}\omega, \tau)$.

(c) $\mathbf{F}$ *is a* π*-tree on* X iff
there exist an isomorphism (φ, ψ) *between* $\mathbf{F}$ *and* $\mathbf{S}$ *and a topology* τ *on* ${}^{\omega}\omega$ *such that*
➢ ψ *is a homeomorphism from* X *onto* $({}^{\omega}\omega, \tau)$ *and*
➢ $\mathbf{S}$ *is a* π*-tree on* $({}^{\omega}\omega, \tau)$.

Proof. (a) Suppose that $\mathbf{F}$ is a locally strict foliage ω,ω-tree with strict branches. Let φ be an order isomorphism from $\mathsf{skeleton}\,\mathbf{F}$ onto the tree $({}^{<\omega}\omega, \subset) = \mathsf{skeleton}\,\mathbf{S}$. For each $p \in {}^{\omega}\omega$, the set $\{x \in {}^{<\omega}\omega : x \subseteq p\}$ is a branch in $\mathbf{S}$, so since $\mathbf{F}$ has strict branches it follows that there is a point $\chi(p)$ in $\mathsf{flesh}\,\mathbf{F}$ such that

$$\big\{\chi(p)\big\} = \bigcap\{\mathbf{F}_{\varphi^{-1}(x)} : x \in {}^{<\omega}\omega \text{ and } x \subseteq p\}.$$

Then it is not hard to prove that the function $\chi{:}{}^{\omega}\omega \to \mathsf{flesh}\,\mathbf{F}$ is a bijection and (φ, χ^{-1}) is an isomorphism between $\mathbf{F}$ and $\mathbf{S}$. The $\leftarrow$ direction follows from (b) of Lemma 6.

(b) Suppose that $\mathbf{F}$ is a Baire foliage tree on X. Let (φ, ψ) be an isomorphism between $\mathbf{F}$ and $\mathbf{S}$, which exists by (a). Then ψ is a bijection from X onto ${}^{\omega}\omega$. Put

$$\tau := \big\{\psi[U] : U \text{ is an open subset of } X\big\};$$

clearly, τ is a topology on ${}^{\omega}\omega$ and ψ is a homeomorphism from X onto $({}^{\omega}\omega, \tau)$. It follows that $\mathbf{S}$ is a Baire foliage tree on $({}^{\omega}\omega, \tau)$ because $\mathbf{F}$ is a Baire foliage tree on X. The $\leftarrow$ direction is similar. Part (c) can be proved by the same argument. □

Corollary 10. *Suppose that* $\mathbf{F}$ *is a Baire foliage tree on a space* X *and* $p \in X$.

(a) $\mathbf{F}$ *is nonincreasing,* $\mathsf{flesh}\,\mathbf{F} = \mathbf{F}_{0_{\mathbf{F}}}$, *and* $\mathsf{height}\,\mathbf{F} = \omega$;
(b) $\mathbf{F}_v$ *is closed-and-open in* X *and* $|\mathbf{F}_v| = 2^{\omega}$ *for all* $v \in \mathsf{nodes}\,\mathbf{F}$;
(c) $\mathsf{scope}_{\mathbf{F}}(p)$ *is a branch in* $\mathbf{F}$;

(d) $\forall n{\in}\omega\ \exists! v{\in}\mathsf{scope}_{\mathbf{F}}(p)\ \big[\,\mathsf{height}_{\mathbf{F}}(v) = n\big]$.

Proof. This corollary is a consequence of (b) of Lemma 9 and (c) of Lemma 6. □

Now we introduce terminology that we need to formulate Theorems 14 and 17.

Definition 11. Suppose $\mathbf{F}$ is a foliage tree and X is a space.

- $\mathsf{span}_{\mathbf{F}}(p,U) := \big\{\,\mathsf{height}_{\mathbf{F}}(v) : v \in \mathsf{scope}_{\mathbf{F}}(p)\ \text{ and }\ \mathsf{shoot}_{\mathbf{F}}(v) \gg \{U\}\big\}$;
- $\mathsf{spectrum}_{\mathbf{F}}(X) := \big\{\,\mathsf{span}_{\mathbf{F}}(p,U) : p \in X\ \text{ and }\ U \in \mathsf{nbhds}(p,X)\big\}$.

Example 12. $\mathsf{span}_{\mathbf{S}}(p, \mathbf{S}_{p\restriction n}) = \omega \setminus n$ for all $p \in {}^{\omega}\omega$ and $n \in \omega$.

Lemma 13. *Suppose that* $\mathbf{F}$ *is a foliage tree and* X *is a space.*

(a) $\mathbf{F}$ *grows into* X iff $\varnothing \notin \mathsf{spectrum}_{\mathbf{F}}(X)$.

(b) *If* $\mathbf{F}$ *is a* π*-tree on* X *and* $p \in X$, *then*

(b1) *the family* $\big\{\,\mathsf{span}_{\mathbf{F}}(p,U) : U \in \mathsf{nbhds}(p,X)\big\}$ *has the* FIP,

(b2) $\bigcap\big\{\,\mathsf{span}_{\mathbf{F}}(p,U) : U \in \mathsf{nbhds}(p,X)\big\} = \varnothing$, *and*

(b3) $\mathsf{span}_{\mathbf{F}}(p,U) \in [\omega]^{\omega}$ *for all* $U \in \mathsf{nbhds}(p,X)$.

Proof. Part (a) is trivial.

(b1) We must show that

$$\text{if}\quad \varepsilon \in [\mathsf{nbhds}(p,X)]^{<\omega} \setminus \{\varnothing\},\quad \text{then}\quad \bigcap_{U\in\varepsilon} \mathsf{span}_{\mathbf{F}}(p,U) \neq \varnothing.$$

For each $U \in \varepsilon$, we have $\mathsf{span}_{\mathbf{F}}(p,U) \supseteq \mathsf{span}_{\mathbf{F}}(p,\bigcap\varepsilon)$ because $U \supseteq \bigcap\varepsilon \neq \varnothing$. Therefore

$$\bigcap_{U\in\varepsilon} \mathsf{span}_{\mathbf{F}}(p,U) \supseteq \mathsf{span}_{\mathbf{F}}(p,\bigcap\varepsilon)$$

and it follows from (a) that $\mathsf{span}_{\mathbf{F}}(p,\bigcap\varepsilon) \neq \varnothing$ since $\bigcap\varepsilon \in \mathsf{nbhds}(p,X)$.

(b2) By (c) of Lemma 9, there exist an isomorphism (φ,ψ) between $\mathbf{F}$ and $\mathbf{S}$ and a topology τ on ${}^{\omega}\omega$ such that ψ is a homeomorphism from X onto $({}^{\omega}\omega,\tau)$ and $\mathbf{S}$ is a π-tree on $({}^{\omega}\omega,\tau)$. Put $q := \psi(p)$. For each $U \subseteq X$, we have $\mathsf{span}_{\mathbf{F}}(p,U) = \mathsf{span}_{\mathbf{S}}\big(q,\psi[U]\big)$ and

$$U \in \mathsf{nbhds}(p,X) \ \leftrightarrow\ \psi[U] \in \mathsf{nbhds}\big(q,({}^{\omega}\omega,\tau)\big).$$

Then it is enough to show that

$$\text{the set}\quad M_q := \bigcap\Big\{\,\mathsf{span}_{\mathbf{S}}(q,V) : V \in \mathsf{nbhds}\big(q,({}^{\omega}\omega,\tau)\big)\Big\}\quad \text{is empty.}$$

It follows from Lemma 6 that $\mathbf{S}_{q\restriction n} \in \mathsf{nbhds}\big(q,({}^{\omega}\omega,\tau)\big)$ for all $n \in \omega$, so using Example 12 we have

$$M_q \subseteq \bigcap\big\{\,\mathsf{span}_{\mathbf{S}}(q,\mathbf{S}_{q\restriction n}) : n \in \omega\big\} = \bigcap\{\omega \setminus n : n \in \omega\} = \varnothing.$$

(b3) It follows from (b1)–(b2) that the set $\mathsf{span}_{\mathbf{F}}(p,U)$ is infinite for all $U \in \mathsf{nbhds}(p,X)$, and $\mathsf{span}_{\mathbf{F}}(p,U) \subseteq \omega$ because $\mathsf{height}\,\mathbf{F} = \omega$ by (a) of Corollary 10. □

4 The first theorem

Theorem 14. *Suppose that* $\mathbf{H}(\lambda)$ *is a* π*-tree on a space* X_λ *for every* $\lambda\in\Lambda$, *where* $2\leqslant|\Lambda|\leqslant\omega$. *Suppose also that for each finite nonempty* $I\subseteq\Lambda$,

- ➢ *if* $R_i\in\mathsf{spectrum}_{\mathbf{H}(i)}(X_i)$ *for all* $i\in I$,
- ➢ *then* $\bigcap_{i\in I}R_i$ *is infinite.*

Then the product $\prod_{\lambda\in\Lambda}X_\lambda$ *has a* π*-tree.*

Corollary 15. *Suppose that* $\mathbf{H}(\lambda)$ *is a* π*-tree on a space* X_λ *and* $\mathsf{cofin}\,\omega\gg\mathsf{spectrum}_{\mathbf{H}(\lambda)}(X_\lambda)$ *for all* $\lambda\in\Lambda$, *where* $1\leqslant|\Lambda|\leqslant\omega$. *Suppose also that a space* Y *has a* π*-tree. Then the product* $Y\times\prod_{\lambda\in\Lambda}X_\lambda$ *also has a* π*-tree.*

Proof of Corollary 15. Let $\mathbf{G}$ be a π-tree on Y and $I\subseteq\Lambda$ be finite and nonempty. Now, if $R\in\mathsf{spectrum}_{\mathbf{G}}(Y)$ and $R_i\in\mathsf{spectrum}_{\mathbf{H}(i)}(X_i)$ for every $i\in I$, then $R\in[\omega]^\omega$ by (b3) of Lemma 13 and it follows from (a) of Lemma 13 that $\bigcap_{i\in I}R_i\supseteq\omega\setminus n$ for some $n\in\omega$. Therefore $R\cap\bigcap_{i\in I}R_i$ is infinite. □

Proof of Theorem 14. We may assume that $2\leqslant\Lambda\in\omega\cup\{\omega\}$. By (c) of Lemma 9, for each $n\in\Lambda$, there exist an isomorphism (φ_n,ψ_n) between $\mathbf{H}(n)$ and $\mathbf{S}$ and a topology τ_n on ${}^\omega\omega$ such that ψ_n is a homeomorphism from X_n onto $({}^\omega\omega,\tau_n)$ and $\mathbf{S}$ is a π-tree on $({}^\omega\omega,\tau_n)$. It follows that

$$\mathsf{spectrum}_{\mathbf{H}(n)}(X_n)=\mathsf{spectrum}_{\mathbf{S}}\big(({}^\omega\omega,\tau_n)\big)\qquad\text{for all }\ n\in\Lambda.$$

Now, for every $k\in\Lambda$, we have the following:

$$\text{if}\quad R_i\in\mathsf{spectrum}_{\mathbf{S}}\big(({}^\omega\omega,\tau_i)\big)\ \text{for every}\ i\in k+1,\quad\text{then}\quad\textstyle\bigcap_{i\in k+1}R_i\ \text{is infinite.}\tag{1}$$

And we must prove that the space $\prod_{n\in\Lambda}({}^\omega\omega,\tau_n)$ has a π-tree.

In this proof we use several specific notations. First, $E\cdot F:=\{e\cup f:e\in E,f\in F\}$. We use this operation in situations when $E\subseteq{}^AC$ and $F\subseteq{}^BC$ with $A\cap B=\varnothing$, so that

$$E\cdot F\ =\ \big\{p\in{}^{A\cup B}C:p{\restriction}A\in E\ \text{ and }\ p{\restriction}B\in F\big\};$$

in particular, when $B=\varnothing$, we have $E\cdot{}^\varnothing C=E$ because ${}^\varnothing C=\{\varnothing\}$. Recall that

$$\textstyle\prod_{i\in I}D_i\ :=\ \big\{\langle p_i\rangle_{i\in I}\in{}^I(\bigcup_{i\in I}D_i)\ :\ p_i\in D_i\ \text{ for all }\ i\in I\big\}.$$

When $v\in{}^{<\omega}\omega$ and $m\in\omega$, we put

$$\widetilde{\mathbf{S}}^m_v\ :=\ \bigcup\{\mathbf{S}_{v^\frown\langle l\rangle}:l\in\omega\setminus m\}.\tag{2}$$

Note that $\{\widetilde{\mathbf{S}}^m_v:m\in\omega\}\gg\mathsf{shoots}_{\mathbf{S}}(v)$ for all $v\in{}^{<\omega}\omega$.

We build a π-tree on the space $\prod_{n\in\Lambda}({}^\omega\omega,\tau_n)=\big({}^\Lambda({}^\omega\omega),\tau\big)$, where τ is the Tychonoff product topology, by using Lemma 16. This lemma states that there exists an indexed family

$$\big\langle\,a(n,v,i)\ :\ n\in\omega,\ v\in{}^{2n}\omega,\ i\in\Lambda\cap(n{+}1)\,\big\rangle$$

such that

(a1) $\forall n\in\omega\ \forall v\in{}^{2n}\omega\ \forall i\in\Lambda\cap(n{+}1)\ \big[a(n,v,i)\in{}^n\omega\big]$;

(a2) $\forall n\in\omega\ \forall v\in{}^{2n}\omega\ \forall m\in\omega$

$$\Big(\big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^m_{a(n,v,i)}\big)\setminus\big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^{m+1}_{a(n,v,i)}\big)\Big)\cdot{}^{\Lambda\cap\{n+1\}}({}^\omega\omega)\ \equiv\ \bigsqcup_{l\in\omega}\prod_{i\in\Lambda\cap(n+2)}\mathbf{S}_{a(n+1,v^\frown\langle m,l\rangle,i)}.$$

Let $\mathbf{G}(\Lambda)$ be a foliage tree with $\mathsf{skeleton}\,\mathbf{G}(\Lambda):=({}^{<\omega}\omega,\subset)$ and with leaves defined as follows:

(b1) $\forall n\in\omega\ \ \forall v\in{}^{2n}\omega$

$$\mathbf{G}(\Lambda)_v \ :=\ \Big(\prod_{i\in\Lambda\cap(n+1)}\mathbf{S}_{a(n,v,i)}\Big)\cdot{}^{\Lambda\smallsetminus(n+1)}({}^{\omega}\omega);$$

(b2) $\forall n\in\omega\ \ \forall v\in{}^{2n}\omega\ \ \forall m\in\omega$

$$\mathbf{G}(\Lambda)_{v^\frown\langle m\rangle} \ :=\ \Big(\Big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^m_{a(n,v,i)}\Big)\smallsetminus\Big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^{m+1}_{a(n,v,i)}\Big)\Big)\cdot{}^{\Lambda\smallsetminus(n+1)}({}^{\omega}\omega).$$

Notice that the construction of $\mathbf{G}(\Lambda)$ doesn't depend on topologies $\tau_n,\ n\in\Lambda$; it depends only on the cardinality of Λ.

To complete the proof, we show that $\mathbf{G}(\Lambda)$ is indeed a π-tree on $\big({}^{\Lambda}({}^{\omega}\omega),\tau\big)$:

☛ $\mathbf{G}(\Lambda)$ is a foliage ω,ω-tree.

☛ $\mathbf{G}(\Lambda)_{0_{\mathbf{G}(\Lambda)}}={}^{\Lambda}({}^{\omega}\omega)$.

We have $0_{\mathbf{G}(\Lambda)}=\langle\rangle$, clause (b1) with $n=0$ says that

$$\mathbf{G}(\Lambda)_{\langle\rangle}\ =\ {}^{\{0\}}\mathbf{S}_{a(0,\langle\rangle,0)}\cdot{}^{\Lambda\smallsetminus 1}({}^{\omega}\omega),$$

so using (8) (see the proof of Lemma 16) we have

$$\mathbf{G}(\Lambda)_{0_{\mathbf{G}(\Lambda)}}\ =\ {}^{\{0\}}\mathbf{S}_{\langle\rangle}\cdot{}^{\Lambda\smallsetminus 1}({}^{\omega}\omega)\ =\ {}^{\{0\}}({}^{\omega}\omega)\cdot{}^{\Lambda\smallsetminus\{0\}}({}^{\omega}\omega)\ =\ {}^{\Lambda}({}^{\omega}\omega).$$

☛ $\mathbf{G}(\Lambda)$ is open in $\big({}^{\Lambda}({}^{\omega}\omega),\tau\big)$.

By (b) of Corollary 10, every set $\mathbf{S}_v$ is closed-and-open in each of spaces $({}^{\omega}\omega,\tau_n)$, and the formula

$$\widetilde{\mathbf{S}}^m_v\ =\ \mathbf{S}_v\smallsetminus\bigcup\{\mathbf{S}_{v^\frown\langle l\rangle}:l\in m\}$$

(which follows from (2)) implies that every set $\widetilde{\mathbf{S}}^m_v$ is closed-and-open in each of $({}^{\omega}\omega,\tau_n)$ too. Therefore every leaf of $\mathbf{G}(\Lambda)$ is open in $\big({}^{\Lambda}({}^{\omega}\omega),\tau\big)$.

☛ $\mathbf{G}(\Lambda)$ is locally strict.

Let $t\in\mathsf{nodes}\,\mathbf{G}(\Lambda)$. First, suppose that $t\in{}^{2n}\omega$ for some $n\in\omega$. Since $\mathbf{S}_u=\widetilde{\mathbf{S}}^0_u$ for all $u\in{}^{<\omega}\omega$, then by (b1) we have

$$\mathbf{G}(\Lambda)_t\ =\ \Big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^0_{a(n,t,i)}\Big)\cdot{}^{\Lambda\smallsetminus(n+1)}({}^{\omega}\omega).$$

Note that for each $u\in{}^{<\omega}\omega$, the $\langle\widetilde{\mathbf{S}}^m_u\rangle_{m\in\omega}$ is a strictly decreasing sequence of sets and $\bigcap_{m\in\omega}\widetilde{\mathbf{S}}^m_u=\varnothing$. Then it follows from (b2) that

$$\mathbf{G}(\Lambda)_t\ \equiv\ \bigsqcup_{m\in\omega}\mathbf{G}(\Lambda)_{t^\frown\langle m\rangle}.$$

Now suppose that $t\in{}^{2n+1}\omega$ for some $n\in\omega$, so that $t=u^\frown\langle m\rangle$ for some $u\in{}^{2n}\omega,\ m\in\omega$. Then by (b2) we have

$$\mathbf{G}(\Lambda)_t\ =\ \mathbf{G}(\Lambda)_{u^\frown\langle m\rangle}\ =\ \Big(\Big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^m_{a(n,u,i)}\Big)\smallsetminus\Big(\prod_{i\in\Lambda\cap(n+1)}\widetilde{\mathbf{S}}^{m+1}_{a(n,u,i)}\Big)\Big)\cdot{}^{\Lambda\cap\{n+1\}}({}^{\omega}\omega)\cdot{}^{\Lambda\smallsetminus(n+2)}({}^{\omega}\omega),$$

so (a2) implies

$$\mathbf{G}(\Lambda)_t\ \equiv\ \bigsqcup_{l\in\omega}\Big(\Big(\prod_{i\in\Lambda\cap(n+2)}\mathbf{S}_{a(n+1,u^\frown\langle m,l\rangle,i)}\Big)\cdot{}^{\Lambda\smallsetminus(n+2)}({}^{\omega}\omega)\Big),$$

and then using (b1) with $v=u^\frown\langle m,l\rangle\in{}^{2(n+1)}\omega$ we have

$$\mathbf{G}(\Lambda)_t\ \equiv\ \bigsqcup_{l\in\omega}\mathbf{G}(\Lambda)_{u^\frown\langle m,l\rangle},\qquad\text{that is,}\qquad\mathbf{G}(\Lambda)_t\ \equiv\ \bigsqcup_{l\in\omega}\mathbf{G}(\Lambda)_{t^\frown\langle l\rangle}.$$

☛ $\mathbf{G}(\Lambda)$ has strict branches.

Suppose that B is a branch in $\mathsf{skeleton}\,\mathbf{G}(\Lambda)$, which means that $B = \{z{\restriction}n : n \in \omega\}$ for some $z \in {}^{\omega}\omega$. Since $\mathbf{G}(\Lambda)$ is a locally strict foliage ω,ω-tree, then $\mathbf{G}(\Lambda)$ is nonincreasing, so

$$\bigcap_{b\in B} \mathbf{G}(\Lambda)_b \;=\; \bigcap_{n\in\omega} \mathbf{G}(\Lambda)_{z{\restriction}2n} \tag{3}$$

because the chain $\{z{\restriction}2n : n \in \omega\}$ is cofinal in $(B,\subset)$. By (b1) we have

$$\mathbf{G}(\Lambda)_{z{\restriction}2n} \;=\; \Big(\prod_{i\in\Lambda\cap(n+1)} \mathbf{S}_{a(n,z{\restriction}2n,i)}\Big)\cdot {}^{\Lambda\setminus(n+1)}({}^{\omega}\omega) \qquad \text{for all } n \in \omega. \tag{4}$$

Since $\mathbf{G}(\Lambda)$ is nonincreasing, it follows from (4) and (a1) that

$$\mathbf{S}_{a(n,z{\restriction}2n,i)} \supset \mathbf{S}_{a(n+1,z{\restriction}2(n+1),i)} \qquad \text{for all } n \in \omega \text{ and } i \in \Lambda \cap (n+1)$$

— that is, for all $i \in \Lambda$ and $n \in \omega \setminus i$. This implies

$$a(n, z{\restriction}2n, i) \subset a\big(n{+}1, z{\restriction}2(n{+}1), i\big) \qquad \text{for all } i \in \Lambda \text{ and } n \in \omega \setminus i,$$

and then, for every $i \in \Lambda$, there is $y_i \in {}^{\omega}\omega$ such that $a(n, z{\restriction}2n, i) \subset y_i$ for all $n \in \omega \setminus i$. Then

$$\bigcap_{n\in\omega\setminus i} \mathbf{S}_{a(n,z{\restriction}2n,i)} = \{y_i\} \qquad \text{for all } i \in \Lambda. \tag{5}$$

Put $y := \langle y_i\rangle_{i\in\Lambda} \in {}^{\Lambda}({}^{\omega}\omega)$. Now (4) and (5) imply $\bigcap_{n\in\omega} \mathbf{G}(\Lambda)_{z{\restriction}2n} = \{y\}$, so the $\bigcap_{b\in B} \mathbf{G}(\Lambda)_b$ is a singleton by (3).

☛ $\mathbf{G}(\Lambda)$ grows into $\big({}^{\Lambda}({}^{\omega}\omega), \tau\big)$.

Suppose that

$$p = \langle p_i\rangle_{i\in\Lambda} \in {}^{\Lambda}({}^{\omega}\omega) \qquad \text{and} \qquad U \in \mathsf{nbhds}\Big(p, \big({}^{\Lambda}({}^{\omega}\omega), \tau\big)\Big).$$

We may assume that

$$U \;=\; \Big(\prod_{i\in k+1} U_i\Big)\cdot {}^{\Lambda\setminus(k+1)}({}^{\omega}\omega) \tag{6}$$

for some $k \in \Lambda$ and some $U_i \in \mathsf{nbhds}\big(p_i, ({}^{\omega}\omega, \tau_i)\big)$ for every $i \in k+1$. Put $R_i := \mathsf{span}_{\mathbf{S}}(p_i, U_i)$ for every $i \in k+1$. Then $\bigcap_{i\in k+1} R_i$ is infinite by (1), so there is some $\bar{n} \in \bigcap_{i\in k+1} R_i$ such that $\bar{n} \geqslant k$. By definition of $\mathsf{span}_{\mathbf{S}}(p_i, U_i)$, for each $i \in k+1$, there is $v_i \in \mathsf{scope}_{\mathbf{S}}(p_i)$ such that

$$\mathsf{height}_{\mathbf{S}}(v_i) = \bar{n} \qquad \text{and} \qquad \mathsf{shoot}_{\mathbf{S}}(v_i) \gg \{U_i\}.$$

This means that for each $i \in k+1$, we have $p_i \in \mathbf{S}_{v_i}$, $v_i \in {}^{\bar{n}}\omega$ (hence $v_i = p_i{\restriction}\bar{n}$), and $\widetilde{\mathbf{S}}^{m_i}_{v_i} = \widetilde{\mathbf{S}}^{m_i}_{p_i{\restriction}\bar{n}} \subseteq U_i$ for some $m_i \in \omega$. Let $\bar{m}$ be the maximal element of $\{m_i : i \in k+1\}$. Then $\widetilde{\mathbf{S}}^{\bar{m}}_{p_i{\restriction}\bar{n}} \subseteq U_i$ for all $i \in k+1$, and hence

$$\prod_{i\in k+1} \widetilde{\mathbf{S}}^{\bar{m}}_{p_i{\restriction}\bar{n}} \;\subseteq\; \prod_{i\in k+1} U_i.$$

Note that $k+1 = \Lambda \cap (k+1)$ because $k \in \Lambda$, so using (6) we get

$$\Big(\prod_{i\in\Lambda\cap(k+1)} \widetilde{\mathbf{S}}^{\bar{m}}_{p_i{\restriction}\bar{n}}\Big)\cdot {}^{\Lambda\setminus(k+1)}({}^{\omega}\omega) \;\subseteq\; U. \tag{7}$$

We already know that $\mathbf{G}(\Lambda)$ is a Baire foliage tree on $\big({}^{\Lambda}({}^{\omega}\omega), \tau\big)$, so using (d) of Corollary 10 we can take node $\bar{v} \in \mathsf{scope}_{\mathbf{G}(\Lambda)}(p)$ such that $\bar{v} \in {}^{2\bar{n}}\omega$. Then, using (b1) with $n = \bar{n}$ and $v = \bar{v}$, we have

$$p \;=\; \langle p_i\rangle_{i\in\Lambda} \;\in\; \mathbf{G}(\Lambda)_{\bar{v}} \;=\; \Big(\prod_{i\in\Lambda\cap(\bar{n}+1)} \mathbf{S}_{a(\bar{n},\bar{v},i)}\Big)\cdot {}^{\Lambda\setminus(\bar{n}+1)}({}^{\omega}\omega),$$

so $p_i \in \mathbf{S}_{a(\bar{n},\bar{v},i)}$ for all $i \in \Lambda\cap(\bar{n}{+}1)$, and hence using (a1) we get $a(\bar{n},\bar{v},i) = p_i{\restriction}\bar{n}$ for all $i \in \Lambda\cap(\bar{n}{+}1)$. Using (b2) and inequality $\bar{n} \geqslant k$ we can write

$$\mathbf{G}(\Lambda)_{\bar{v}^\frown\langle m\rangle} \subseteq \Big(\prod_{i\in\Lambda\cap(\bar{n}+1)} \widetilde{\mathbf{S}}^m_{p_i{\restriction}\bar{n}}\Big)\cdot {}^{\Lambda\setminus(\bar{n}+1)}({}^\omega\omega) \subseteq \Big(\prod_{i\in\Lambda\cap(k+1)} \widetilde{\mathbf{S}}^m_{p_i{\restriction}\bar{n}}\Big)\cdot {}^{\Lambda\setminus(k+1)}({}^\omega\omega) \qquad \text{for all } m\in\omega.$$

Therefore using (7) we get $\mathbf{G}(\Lambda)_{\bar{v}^\frown\langle m\rangle} \subseteq U$ for all $m \in \omega\setminus\bar{m}$. This means that we have found $\bar{v} \in \mathsf{scope}_{\mathbf{G}(\Lambda)}(p)$ such that $\mathsf{shoot}_{\mathbf{G}(\Lambda)}(\bar{v}) \gg \{U\}$. □

Lemma 16. *For each $\Lambda \in \big(\omega\cup\{\omega\}\big)\setminus 2$, there is a family $\big\langle a(n,v,i) : n\in\omega,\ v\in{}^{2n}\omega,\ i\in\Lambda\cap(n{+}1)\big\rangle$ such that*

(a1) $\forall n{\in}\omega\ \forall v{\in}{}^{2n}\omega\ \forall i{\in}\Lambda\cap(n{+}1)\ \big[a(n,v,i)\in{}^n\omega\big]$;

(a2) $\forall n{\in}\omega\ \forall v{\in}{}^{2n}\omega\ \forall m{\in}\omega$

$$\Big(\big(\prod_{i\in\Lambda\cap(n+1)} \widetilde{\mathbf{S}}^m_{a(n,v,i)}\big)\setminus\big(\prod_{i\in\Lambda\cap(n+1)} \widetilde{\mathbf{S}}^{m+1}_{a(n,v,i)}\big)\Big)\cdot {}^{\Lambda\cap\{n+1\}}({}^\omega\omega) \equiv \bigsqcup_{l\in\omega}\prod_{i\in\Lambda\cap(n+2)} \mathbf{S}_{a(n+1,v^\frown\langle m,l\rangle,i)}.$$

Proof. We construct this indexed family by recursion on $n\in\omega$ as follows:

When $n=0$, we have ${}^{2n}\omega = {}^n\omega = {}^0\omega = \big\{\langle\rangle\big\}$ and $\Lambda\cap(n+1) = \{0\}$ because $\Lambda\geqslant 2$, so (a1) with $n=0$ just says

$$a\big(0,\langle\rangle,0\big) = \langle\rangle. \tag{8}$$

When $n=1$, we must choose $a(1,v,i)\in{}^1\omega$ (for all $v\in{}^2\omega$ and $i\in\Lambda\cap 2$) in such a way that (a2) with $n=0$ is satisfied. Since $\Lambda\geqslant 2$, then $\Lambda\cap 1=\{0\}$ and $\Lambda\cap 2=\{0,1\}$, so (a2) with $n=0$ says that

$$\big({}^{\{0\}}\widetilde{\mathbf{S}}^m_{a(0,\langle\rangle,0)}\setminus{}^{\{0\}}\widetilde{\mathbf{S}}^{m+1}_{a(0,\langle\rangle,0)}\big)\cdot{}^{\{1\}}({}^\omega\omega) \equiv \bigsqcup_{l\in\omega}\prod_{i\in\{0,1\}}\mathbf{S}_{a(1,\langle\rangle^\frown\langle m,l\rangle,i)} \qquad \text{for all } m\in\omega.$$

Using (2) and (8), this can be simplified to

$${}^{\{0\}}\mathbf{S}_{\langle m\rangle}\cdot{}^{\{1\}}({}^\omega\omega) \equiv \bigsqcup_{l\in\omega}\prod_{i\in\{0,1\}}\mathbf{S}_{a(1,\langle m,l\rangle,i)} \qquad \text{for all } \forall m\in\omega.$$

Then we can take $a\big(1,\langle m,l\rangle,0\big) := \langle m\rangle$ and $a\big(1,\langle m,l\rangle,1\big) := \langle l\rangle$ for every $m,l\in\omega$.

When $n\geqslant 2$, the choice of $a(n,v,i)$ can be carried out similar to the case $n=1$ if we note that

$${}^\omega\omega \equiv \bigsqcup_{a\in{}^h\omega}\mathbf{S}_a \qquad \text{for all } h\in\omega,$$

and that for every $k\geqslant 2$, every $a=\langle a_i\rangle_{i\in k}\in{}^k({}^{2n}\omega)$, and every $m\in\omega$,

$$\big(\prod_{i\in k}\widetilde{\mathbf{S}}^m_{a_i}\big)\setminus\big(\prod_{i\in k}\widetilde{\mathbf{S}}^{m+1}_{a_i}\big) = \big(\prod_{i\in k}\bigcup_{l\in\omega\setminus m}\mathbf{S}_{a_i^\frown\langle l\rangle}\big)\setminus\big(\prod_{i\in k}\bigcup_{l\in\omega\setminus(m+1)}\mathbf{S}_{a_i^\frown\langle l\rangle}\big) =$$

$$= \bigcup\Big\{\prod_{i\in k}\mathbf{S}_{a_i^\frown\langle l_i\rangle} : l=\langle l_i\rangle_{i\in k}\in{}^k(\omega\setminus m)\Big\}\setminus\bigcup\Big\{\prod_{i\in k}\mathbf{S}_{a_i^\frown\langle l_i\rangle} : l=\langle l_i\rangle_{i\in k}\in{}^k\big(\omega\setminus(m+1)\big)\Big\} \equiv$$

$$\equiv \bigsqcup\Big\{\prod_{i\in k}\mathbf{S}_{a_i^\frown\langle l_i\rangle} : l=\langle l_i\rangle_{i\in k}\in{}^k(\omega\setminus m)\setminus{}^k\big(\omega\setminus(m{+}1)\big)\Big\}$$

and the set ${}^k(\omega\setminus m)\setminus{}^k\big(\omega\setminus(m{+}1)\big)$ is infinite. □

5 The second theorem

Theorem 17.

(a) *Suppose that $\mathbf{F}(\alpha)$ is a π-tree on a space X_α for every $\alpha\in A$, where $1\leqslant|A|\leqslant\omega$. Suppose also that for each $\alpha\in A$, there is $\gamma_\alpha\subseteq\mathsf{power.set}(\omega)$ such that*

➢ $|\gamma_\alpha| \leqslant \omega$,

➢ γ_α *has the* FIP*, and*

➢ $\gamma_\alpha \gg \mathsf{spectrum}_{\mathbf{F}(\alpha)}(X_\alpha)$.

Then the product $\prod_{\alpha\in A} X_\alpha$ *has a* π*-tree.*

(b) *Suppose, in addition to* (a)*, that* $\mathbf{G}$ *is a* π*-tree on a space* Y *and* $\mathsf{spectrum}_{\mathbf{G}}(Y)$ *has the* FIP*. Then the product* $Y \times \prod_{\alpha\in A} X_\alpha$ *also has a* π*-tree.*

Lemma 18. *Suppose that* $2 \leqslant \Lambda \in \omega\cup\{\omega\}$ *and for each* $n \in \Lambda$, $\varnothing \neq \delta_n \subseteq \mathsf{power.set}(\omega)\setminus\{\varnothing\}$ *and* $\bigcap \delta_n = \varnothing$*. Suppose also that* δ_0 *has the* FIP *and for each* $n \in \Lambda\setminus\{0\}$*, there is* $\gamma_n \subseteq \mathsf{power.set}(\omega)$ *such that*

➢ $|\gamma_n| \leqslant \omega$,

➢ γ_n *has the* FIP*, and*

➢ $\gamma_n \gg \delta_n$.

Then there exists a sequence $\langle\alpha_n\rangle_{n\in\Lambda}$ *of strictly increasing functions* $\alpha_n{:}\omega\to\omega$ *such that*

(♣) *if* $k \in \Lambda$ *and* $A_i \in \{\alpha_i[D] : D \in \delta_i\}$ *for all* $i \leqslant k$*, then* $\bigcap_{i\leqslant k} A_i$ *is infinite.*

Lemma 19. *Suppose that* $\mathbf{F}$ *is a* π*-tree on a space* X *and* $\alpha{:}\omega\to\omega$ *is a strictly increasing function. Then there exists a* π*-tree* $\mathbf{H}$ *on* X *such that*

(♥) $\alpha\big[\mathsf{span}_{\mathbf{F}}(p,U)\big] \subseteq \mathsf{span}_{\mathbf{H}}(p,U)$ *for all* $p \in X$ *and* $U \in \mathsf{nbhds}(p,X)$.

Proof of Theorem 17. Note that part (a) follows from part (b). Indeed, let $\beta \in A$, $\mathbf{G} := \mathbf{F}(\beta)$, and $Y := X_\beta$. Since γ_β has the FIP, $\gamma_\beta \gg \mathsf{spectrum}_{\mathbf{G}}(Y)$, and (by (a) of Lemma 13) $\varnothing \notin \mathsf{spectrum}_{\mathbf{G}}(Y)$, then $\mathsf{spectrum}_{\mathbf{G}}(Y)$ also has the FIP. The case when $|A| = 1$ is trivial, so we may assume that $A\setminus\{\beta\} \neq \varnothing$, and then the space

$$\prod_{\alpha\in A} X_\alpha \;=\; Y \times \prod_{\alpha\in A\setminus\{\beta\}} X_\alpha$$

has a π-tree by (b).

To prove (b) it is convenient to assume that $A = \Lambda\setminus\{0\}$ and $2 \leqslant \Lambda \in \omega\cup\{\omega\}$. Put $X_0 := Y$ and $\mathbf{F}(0) := \mathbf{G}$; then we must prove that the space $\prod_{n\in\Lambda} X_n$ has a π-tree. Let $\delta_n := \mathsf{spectrum}_{\mathbf{F}(n)}(X_n)$ for every $n \in \Lambda$. Then using Lemma 13 we see that $\delta_n \subseteq \mathsf{power.set}(\omega)\setminus\{\varnothing\}$ and $\bigcap\delta_n = \varnothing$ for all $n \in \Lambda$, so we can apply Lemma 18. Then we get a sequence $\langle\alpha_n\rangle_{n\in\Lambda}$ of strictly increasing functions $\alpha_n{:}\omega\to\omega$ such that condition (♣) holds. Next, applying Lemma 19 to $\mathbf{F}_n$, X_n, and α_n for every $n \in \Lambda$, we obtain a sequence $\big\langle\mathbf{H}(n)\big\rangle_{n\in\Lambda}$ such that for every $n \in \Lambda$, $\mathbf{H}(n)$ is a π-tree on X_n and

$$\alpha_n\big[\mathsf{span}_{\mathbf{F}(n)}(p,U)\big] \subseteq \mathsf{span}_{\mathbf{H}(n)}(p,U) \qquad \text{for all } p \in X_n \text{ and } U \in \mathsf{nbhds}(p,X_n). \tag{9}$$

Now we can use Theorem 14 to show that the product $\prod_{n\in\Lambda} X_n$ has a π-tree. Suppose that $I \subseteq \Lambda$ is finite and nonempty; then $I \subseteq k+1$ for some $k \in \Lambda$. Let $R_i \in \mathsf{spectrum}_{\mathbf{H}(i)}(X_i)$ for every $i \in k+1$; we must show that the set $\bigcap_{i\in I} R_i$ is infinite. For each $i \in k+1$, $R_i = \mathsf{span}_{\mathbf{H}(i)}(p_i,U_i)$ for some $p_i \in X_i$ and $U_i \in \mathsf{nbhds}(p_i,X_i)$. Put $A_i := \alpha_i\big[\mathsf{span}_{\mathbf{F}(i)}(p_i,U_i)\big]$ for every $i \in k+1$; then by (9) we have $A_i \subseteq R_i$. Now,

$$A_i \in \big\{\alpha_i[R] : R \in \mathsf{spectrum}_{\mathbf{F}(i)}(X_i)\big\} = \big\{\alpha_i[D] : D \in \delta_i\big\} \qquad \text{for all } i \in k+1,$$

so (by (♣) of Lemma 18) the $\bigcap_{i\in k+1} A_i$ is infinite, hence the $\bigcap_{i\in k+1} R_i$ is infinite, and then the $\bigcap_{i\in I} R_i$ is infinite too. □

Proof of Lemma 18. It is not hard to show that each γ_n is not empty, so we may assume that $\gamma_n = \{G_i(n) : i \in \omega\}$ for every $n \in \Lambda\setminus\{0\}$. Since δ_0 has the FIP and $\bigcap\delta_0 = \varnothing$, then

$$\bigcap\varepsilon \text{ is infinite} \qquad \text{for all } \varepsilon \in [\delta_0]^{<\omega}\setminus\{\varnothing\}. \tag{10}$$

Also $\bigcap\gamma_n = \varnothing$ for all $n \in \Lambda \setminus \{0\}$ (because $\gamma_n \gg \delta_n$, $\varnothing \notin \delta_n \neq \varnothing$, and $\bigcap\delta_n = \varnothing$), so by the same reasons we get

$$\bigcap_{j\leqslant i} G_j(n) \text{ is infinite} \qquad \text{for all } n \in \Lambda \setminus \{0\} \text{ and } i \in \omega. \tag{11}$$

Now, using (11), for every $n \in \Lambda \setminus \{0\}$ and $i \in \omega$, we can choose $f_i(n) \in \bigcap_{j\leqslant i} G_j(n)$ in such a way that

$$f_{i+1}(n) > f_i(n) \qquad \text{for all } n \in \Lambda \setminus \{0\} \text{ and } \forall i \in \omega. \tag{12}$$

Put $F(n) := \{f_i(n) : i \in \omega\}$ for every $n \in \Lambda \setminus \{0\}$; then $\{F(n) \setminus m : m \in \omega\} \gg \gamma_n$ for all $n \in \Lambda \setminus \{0\}$, and hence

$$\{F(n) \setminus m : m \in \omega\} \gg \delta_n \qquad \text{for all } n \in \Lambda \setminus \{0\}. \tag{13}$$

Let $F(0) \in \delta_0$; then $F(0)$ is infinite by (10), so we may assume that $F(0) = \{f_i(0) : i \in \omega\}$ and $f_{i+1}(0) > f_i(0)$ for all $i \in \omega$. Put $h_{-1} := -1$ and $f_{-1}(n) := -1$ for every $n \in \Lambda$. By recursion on $i \in \omega$, we can build a strictly increasing sequence $\langle h_i\rangle_{i\in\omega}$ of natural numbers in such a way that

$$h_i \;>\; h_{i-1} + f_{i-j}(j) - f_{i-j-1}(j) \qquad \text{for all } i \in \omega \text{ and } j \in \Lambda \cap (i+1). \tag{14}$$

Let $\langle\beta_n\rangle_{n\in\Lambda}$ be a sequence of functions with

$$\mathsf{domain}\,\beta_n \;:=\; F(n) \cup \{-1\} \;=\; \{f_l(n) : l \in \omega \cup \{-1\}\}$$

and such that

$$\beta_n\big(f_l(n)\big) = h_{n+l} \qquad \text{for all } n \in \Lambda \text{ and } l \in \omega \cup \{-1\}. \tag{15}$$

Note that (15) implies

$$\beta_n\big[F(n)\big] \;=\; \{h_j : j \in \omega \setminus n\} \qquad \text{for all } n \in \Lambda. \tag{16}$$

Now, for all $n \in \Lambda$ and $l \in \omega$, (14) with $i = n + l$, $j = n$ says that

$$h_{n+l} - h_{n+l-1} \;>\; f_{n+l-n}(n) - f_{n+l-n-1}(n) \;=\; f_l(n) - f_{l-1}(n),$$

so by (15) we have

$$\beta_n\big(f_l(n)\big) - \beta_n\big(f_{l-1}(n)\big) \;>\; f_l(n) - f_{l-1}(n) \qquad \text{for all } n \in \Lambda \text{ and } l \in \omega.$$

This means that for each $n \in \Lambda$, we can choose a strictly increasing function $\alpha_n{:}\omega \to \omega$ such that $\alpha_n{\restriction}F(n) = \beta_n{\restriction}F(n)$.

Now we prove that condition (♣) is satisfied. Suppose that $k \in \Lambda$ and for every $i \in k+1$, $A_i = \alpha_i\big[D(i)\big]$ for some $D(i) \in \delta_i$. Using (13), for each $i \in (k+1) \setminus \{0\}$, we can choose some $m_i \in \omega$ such that $D(i) \supseteq F(i) \setminus m_i$. Therefore, by (16), for each $i \in (k+1) \setminus \{0\}$, we can find some $l_i \in \omega$ such that $A_i \supseteq \{h_j : j \in \omega \setminus l_i\}$. It follows that

$$\bigcap_{i\in(k+1)\setminus\{0\}} A_i \;\supseteq\; \{h_j : j \in \omega \setminus l\} \qquad \text{for some } l \in \omega.$$

Now, $D(0) \cap F(0)$ is infinite by (10), $\alpha_0\big[F(0)\big] = \{h_j : j \in \omega\}$ by (16), and α_0 is injective, therefore $A_0 \cap \{h_j : j \in \omega\}$ is infinite. This means that $\bigcap_{i\in k+1} A_i$ is infinite too. □

Proof of Lemma 19. In this proof we apply the foliage hybrid operation, see details in Section 8. Put $\alpha(-1) := -1$. Suppose that $v \in \mathsf{nodes}\,\mathbf{F}$. Set

$$k(v) \;:=\; \alpha\big(\mathsf{height}_{\mathbf{F}}(v)\big) - \alpha\big(\mathsf{height}_{\mathbf{F}}(v) - 1\big);$$

then $k(v) \in \omega \setminus \{0\}$. Let $\mathcal{T}(v)$ be a tree isomorphic to the tree $({}^{<k(v)+1}\omega, \subset)$ and such that $0_{\mathcal{T}(v)} = v$ and $\max\mathcal{T}(v) = \mathsf{sons}_{\mathbf{F}}(v)$. Let $\mathbf{G}(v)$ be a foliage tree with $\mathsf{skeleton}\,\mathbf{G}(v) := \mathcal{T}(v)$ and with leaves defined by recursion on $i \in k(v)+1$ as follows:

(base) If $i = 0$ and $t \in \mathsf{level}_{\mathcal{T}(v)}\big(k(v) - i\big)$ $\big($that is, if $t \in \mathsf{max}\,\mathcal{T}(v)\,\big)$,
then $\mathbf{G}(v)_t := \mathbf{F}_t$.

(step) If $1 \leqslant i \leqslant k(v)$ and $t \in \mathsf{level}_{\mathcal{T}(v)}\big(k(v) - i\big)$,
then $\mathbf{G}(v)_t := \bigcup\big\{\mathbf{G}(v)_s : s \in \mathsf{sons}_{\mathcal{T}(v)}(t)\big\}$.

It is not hard to show the following (we use here the terminology of Definition 29):

(c1) $0_{\mathbf{G}(v)} = v$ and $\mathsf{max}\,\mathbf{G}(v) = \mathsf{sons}_{\mathbf{F}}(v) = \mathsf{level}_{\mathbf{G}(v)}\big(k(v)\big)$;

(c2) $\mathbf{G}(v)_v = \mathbf{F}_v$;

(c3) $\mathbf{G}(v)$ is a foliage graft for $\mathbf{F}$;

(c4) $\mathsf{cut}\big(\mathbf{F}, \mathbf{G}(v)\big) = \varnothing$;

(c5) $\mathbf{G}(v)$ is ω-branching, locally strict, open in X, and has bounded chains;

(c6) $\mathsf{height}\,\mathbf{G}(v) = k(v) + 1$;

(c7) $\mathsf{shoot}_{\mathbf{G}(v)}(t) \gg \mathsf{shoot}_{\mathbf{F}}(v)$ for all $t \in \mathsf{nodes}\,\mathbf{G}(v) \setminus \mathsf{max}\,\mathbf{G}(v)$;

(c8) $\mathsf{explant}\big(\mathbf{F}, \mathbf{G}(v)\big) = \varnothing$.

Now let $\varphi := \big\{\mathbf{G}(v) : v \in \mathsf{nodes}\,\mathbf{F}\big\}$. We may assume that

$$\mathsf{implant}\,\mathbf{G}(v) \cap \mathsf{implant}\,\mathbf{G}(u) = \varnothing \qquad \text{for all } v \neq u \in \mathsf{nodes}\,\mathbf{F},$$

so φ is a consistent family of foliage grafts for $\mathbf{F}$. Let $\mathbf{H} := \mathsf{fol.hybr}(\mathbf{F}, \varphi)$; note that $\mathsf{loss}(\mathbf{F}, \varphi) = \varnothing$ by (c4). By induction on $\mathsf{height}_{\mathbf{F}}(v)$, we can prove that

$$\mathsf{height}_{\mathbf{H}}(v) \;=\; \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v) - 1\big) + 1 \qquad \text{for all } v \in \mathsf{nodes}\,\mathbf{F}. \tag{17}$$

Indeed, if $\mathsf{height}_{\mathbf{F}}(v) = 0$, then $v = 0_{\mathbf{F}}$, so $v = 0_{\mathbf{H}}$, and hence

$$\mathsf{height}_{\mathbf{H}}(v) \;=\; 0 \;=\; \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v) - 1\big) + 1.$$

If $\mathsf{height}_{\mathbf{F}}(v) \geqslant 1$, then let t be the node in $\mathbf{F}$ such that $v \in \mathsf{sons}_{\mathbf{F}}(t)$, and then inductively we can write

$$\begin{gathered}\mathsf{height}_{\mathbf{H}}(v) \;=\; \mathsf{height}_{\mathbf{H}}(t) + \mathsf{height}_{\mathbf{G}(t)}(v) \;=\; \mathsf{height}_{\mathbf{H}}(t) + \big(\,\mathsf{height}\,\mathbf{G}(t) - 1\big) \;=\\ \mathsf{height}_{\mathbf{H}}(t) + \big(k(t) + 1\big) - 1 \;=\; \alpha\big(\,\mathsf{height}_{\mathbf{F}}(t) - 1\big) + 1 + k(t) \;=\\ \alpha\big(\,\mathsf{height}_{\mathbf{F}}(t) - 1\big) + 1 + \alpha\big(\,\mathsf{height}_{\mathbf{F}}(t)\big) - \alpha\big(\,\mathsf{height}_{\mathbf{F}}(t) - 1\big) \;=\\ \alpha\big(\,\mathsf{height}_{\mathbf{F}}(t)\big) + 1 \;=\; \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v) - 1\big) + 1.\end{gathered}$$

Now, (c4)–(c6) with Lemma 31 say that $\mathbf{H}$ is a Baire foliage tree on X and (c7)–(c8) imply that each $\mathbf{G}(v)$ preserves shoots of $\mathbf{F}$ (see Definition 32), so $\mathbf{H}$ grows into X by Lemmas 33 and 34. Therefore $\mathbf{H}$ is a π-tree on X.

Let us show that (♥) holds. Suppose that $p \in X$, $U \in \mathsf{nbhds}(p, X)$, and $r \in \mathsf{span}_{\mathbf{F}}(p, U)$. Then $r = \mathsf{height}_{\mathbf{F}}(v)$ for some node $v \in \mathsf{scope}_{\mathbf{F}}(p)$ such that $\mathsf{shoot}_{\mathbf{F}}(v) \gg \{U\}$. Let s be the node in $\mathsf{sons}_{\mathbf{F}}(v)$ such that $p \in \mathbf{F}_s$ and let t be the node in $\mathbf{G}(v)$ such that $s \in \mathsf{sons}_{\mathbf{G}(v)}(t)$. Then $t \in \mathsf{scope}_{\mathbf{H}}(p)$ and using (c7) and (a) of Proposition 30 we obtain

$$\mathsf{shoot}_{\mathbf{H}}(t) = \mathsf{shoot}_{\mathbf{G}(v)}(t) \gg \mathsf{shoot}_{\mathbf{F}}(v), \tag{18}$$

so $\mathsf{shoot}_{\mathbf{H}}(t) \gg \{U\}$, and hence $\mathsf{height}_{\mathbf{H}}(t) \in \mathsf{span}_{\mathbf{H}}(p, U)$. Therefore to complete the proof it is enough to show that $\alpha(r) = \mathsf{height}_{\mathbf{H}}(t)$. Indeed, using (c6) and (17) we have

$$\begin{gathered}\mathsf{height}_{\mathbf{H}}(t) \;=\; \mathsf{height}_{\mathbf{H}}(v) + \mathsf{height}_{\mathbf{G}(v)}(t) \;=\; \mathsf{height}_{\mathbf{H}}(v) + \mathsf{height}_{\mathbf{G}(v)}(s) - 1 \;=\\ \mathsf{height}_{\mathbf{H}}(v) + \big(\,\mathsf{height}\,\mathbf{G}(v) - 1\big) - 1 \;=\; \mathsf{height}_{\mathbf{H}}(v) + \big(k(v) + 1\big) - 2 \;=\\ \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v) - 1\big) + 1 + \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v)\big) - \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v) - 1\big) - 1 \;=\; \alpha\big(\,\mathsf{height}_{\mathbf{F}}(v)\big) \;=\; \alpha(r).\end{gathered}$$

□

6 Nice π-tree for a co-countable subspace

In this section we prove Corollary 21, which states that if a space X has a "very nice" π-tree (that is, a π-tree $\mathbf{F}$ such that $\mathsf{cofin}\,\omega \gg \mathsf{spectrum}_{\mathbf{F}}(X)$) and if $A \subseteq X$ is at most countable, then the subspace $X \setminus A$ has a "nice" π-tree — that is, a π-tree that satisfies the conditions of Theorem 17. This result allows to apply Theorem 17 to co-countable subspaces of the Sorgenfrey line, see (c) of Lemma 24 and Corollary 25 in Section 7.

Proposition 20. *Suppose that $\mathbf{F}$ is a Baire foliage tree on a space X and $A \subseteq X$ is at most countable. Then there exists a Baire foliage tree $\mathbf{H}$ on the subspace $X \setminus A$ such that for every $p \in X \setminus A$, there is a strictly increasing function $f_p\colon \omega \to \omega$ with a property*

($\blacklozenge$) $\big\{ 2n+1 : n \in \omega \text{ and } f_p(n) \in \mathsf{span}_{\mathbf{F}}(p, U) \big\} \subseteq \mathsf{span}_{\mathbf{H}}(p, U \setminus A)$ *for all* $U \in \mathsf{nbhds}(p, X)$.

Corollary 21. *Suppose that $\mathbf{F}$ is a π-tree on a space X such that $\mathsf{cofin}\,\omega \gg \mathsf{spectrum}_{\mathbf{F}}(X)$ and $A \subseteq X$ is at most countable. Then there exists a π-tree $\mathbf{H}$ on the subspace $X \setminus A$ such that*

($\spadesuit$) $\mathsf{cofin}\{2n+1 : n \in \omega\} \gg \mathsf{spectrum}_{\mathbf{H}}(X \setminus A)$.

Remark 22. In statements of Proposition 20 and Corollaries 21 the sequence $\langle 2n+1 \rangle_{n\in\omega}$ can be replaced by an arbitrary sequence $\langle k_n \rangle_{n\in\omega}$ of natural numbers such that $k_0 \geqslant 1$ and $k_{n+1} > k_n + 1$ for all $n \in \omega$.

Proof of Corollary 21. Let $\mathbf{H}$ be a Baire foliage tree on the subspace $X \setminus A$ from Proposition 20. First we show that condition ($\spadesuit$) is satisfied. Suppose that $D \in \mathsf{spectrum}_{\mathbf{H}}(X \setminus A)$; that is, $D = \mathsf{span}_{\mathbf{H}}(p, U \setminus A)$ for some $p \in X \setminus A$ and $U \in \mathsf{nbhds}(p, X)$. Let $f_p\colon \omega \to \omega$ be a function that satisfies condition ($\blacklozenge$) of Proposition 20. Since $\mathbf{F}$ is a π-tree on X, then $\mathsf{span}_{\mathbf{F}}(p, U) \neq \varnothing$ by (a) of Lemma 13, so it follows from $\mathsf{cofin}\,\omega \gg \mathsf{spectrum}_{\mathbf{F}}(X)$ that there is some $\bar{m} \in \omega$ such that $\omega \setminus \bar{m} \subseteq \mathsf{span}_{\mathbf{F}}(p, U)$. Therefore, since f_p is strictly increasing, there is some $\bar{n} \in \omega$ such that $f_p(n) \in \mathsf{span}_{\mathbf{F}}(p, U)$ for all $n \geqslant \bar{n}$. Then by ($\blacklozenge$) we have

$$\big\{ 2n+1 : n \in \omega \setminus \bar{n} \big\} \subseteq \mathsf{span}_{\mathbf{H}}(p, U \setminus A) = D,$$

hence ($\spadesuit$) is satisfied.

It follows from the above reasoning that $D \neq \varnothing$, so $\varnothing \notin \mathsf{spectrum}_{\mathbf{H}}(X \setminus A)$, and hence $\mathbf{H}$ grows into $X \setminus A$ by (a) of Lemma 13. This means that $\mathbf{H}$ is a π-tree on $X \setminus A$. □

In the following lemma we use terminology of the foliage hybrid operation, see Definition 29 in Section 8.

Lemma 23. *Suppose that $\mathbf{F}$ is a Baire foliage tree on a space X, $p \in X$, and $v \in \mathsf{scope}_{\mathbf{F}}(p)$. Then there exists a foliage tree $\mathbf{G}$ such that*

(d1) $0_{\mathbf{G}} = v$ *and* $\mathsf{max}\,\mathbf{G} = \mathsf{sons}_{\mathbf{G}}(0_{\mathbf{G}})$;

(d2) $\mathbf{G}_v = \mathbf{F}_v \setminus \{p\} \equiv \bigsqcup_{m \in \mathsf{max}\,\mathbf{G}} \mathbf{F}_m$;

(d3) $\mathbf{G}$ *is a foliage graft for* $\mathbf{F}$;

(d4) $\mathsf{implant}\,\mathbf{G} = \varnothing$;

(d5) $\mathbf{G}$ *is ω-branching, locally strict, open in X, has bounded chains, and* $\mathsf{height}\,\mathbf{G} = 2$.

Proof of Lemma 23. Put

$$B := \bigcup \big\{ \mathsf{sons}_{\mathbf{F}}(u) : u \in \mathsf{scope}_{\mathbf{F}}(p) \cap v{\downarrow_{\mathbf{F}}} \big\} \qquad \text{and} \qquad \mathsf{MAX} := B \setminus \mathsf{scope}_{\mathbf{F}}(p).$$

Let $\mathcal{T}$ be a partial order such that

$$\mathsf{nodes}\,\mathcal{T} := \{v\} \cup \mathsf{MAX} \qquad \text{and} \qquad <_{\mathcal{T}} := \big\{ (v, m) : m \in \mathsf{MAX} \big\}.$$

Then $\mathcal{T}$ is a tree, $0_{\mathcal{T}} = v$, $\mathsf{max}\,\mathcal{T} = \mathsf{MAX}$, and $\mathcal{T}$ is a graft for $\mathsf{skeleton}\,\mathbf{F}$. Now let $\mathbf{G}$ be a foliage tree with $\mathsf{skeleton}\,\mathbf{G} := \mathcal{T}$ and with leaves $\mathbf{G}_v := \mathbf{F}_v \setminus \{p\}$ and $\mathbf{G}_m := \mathbf{F}_m$ for all $m \in \mathsf{max}\,\mathcal{T}$. Then using (b) of Lemma 9 and Corollary 10 it is not hard to verify that clauses (d1)–(d5) are satisfied. □

Proof of Proposition 20. We may assume that $A = \{p_i : i \in |A|\}$ and $p_i \neq p_j$ for all $i \neq j \in |A|$. First we build sequences $\langle M_i\rangle_{i\in|A|}$, $\langle z_i\rangle_{i\in|A|}$, and $\langle \mathbf{G}(i)\rangle_{i\in|A|}$ by recursion on $i \in |A|$:

(e1) $M_0 := \{0_{\mathbf{F}}\}$;

(e2) $z_i :=$ the node in $\mathbf{F}$ such that $\{z_i\} = M_i \cap \mathsf{scope}_{\mathbf{F}}(p_i)$;

(e3) $\mathbf{G}(i) :=$ the foliage tree $\mathbf{G}$ from Lemma 23 with $p = p_i$ and $v = z_i$;

(e4) $M_{i+1} := \big(M_i \setminus \{z_i\}\big) \cup \bigcup\big\{\mathsf{sons}_{\mathbf{F}}(m) : m \in \mathsf{max}\,\mathbf{G}(i)\big\}$.

The correctness of clause (e2) follows from (f3), see below.

It is not hard to verify that for each $i \in |A|$, the following conditions are satisfied:

(f1) M_i is an antichain in $\mathbf{F}$ (that is, $u \not\leq_{\mathbf{F}} v$ and $v \not\leq_{\mathbf{F}} u$ for all $u, v \in M_i$);

(f2) $M_{i+1} \subseteq (M_i)\!\downarrow_{\mathbf{F}}$;

(f3) $X \setminus \{p_j : j \in i\} \equiv \bigsqcup_{m\in M_i} \mathbf{F}_m$;

(f4) $0_{\mathbf{G}(i)} \in M_i$;

(f5) $\mathsf{cut}\big(\mathbf{F}, \mathbf{G}(i)\big) = \{p_i\}$;

(f6) $\big\{\mathbf{G}(j) : j \in i+1\big\}$ is a consistent family of foliage grafts for $\mathbf{F}$.

Now it follows from (f6) that $\varphi := \big\{\mathbf{G}(i) : i \in |A|\big\}$ is a consistent family of foliage grafts for $\mathbf{F}$, so we can define $\mathbf{H} := \mathsf{fol.hybr}(\mathbf{F}, \varphi)$. Then $\mathbf{H}$ is a Baire foliage tree on $X \setminus A$ by Lemma 31, (f5), and (d5). Also $\mathbf{H}$ satisfies the following:

(g1) $\mathbf{H}$ has nonempty leaves.
This follows from (b) of Corollary 10.

(g2) $\mathsf{nodes}\,\mathbf{H} \subseteq \mathsf{nodes}\,\mathbf{F}$.
This follows from (d4).

(g3) $\mathbf{H}_u = \mathbf{F}_u \setminus A$ for all $u \in \mathsf{nodes}\,\mathbf{H}$.
This also follows from (d4).

(g4) $\mathsf{height}_{\mathbf{H}}(u) \in \{2n : n \in \omega\}$ for all $u \in M_i$ and $i \in |A|$.
We prove (g4) by induction on $i \in |A|$. Obviously, $\mathsf{height}_{\mathbf{H}}(u)$ is even when $u \in M_0$. Assume as inductive hypothesis that the assertion of (g4) holds for all $u \in \bigcup_{i\leqslant k} M_i$ and prove it for an arbitrary $u \in M_{k+1}$. If $u \in M_k$, then $\mathsf{height}_{\mathbf{H}}(u)$ is even by the inductive hypothesis. If $u \notin M_k$, then (e4) implies that $u \in \mathsf{sons}_{\mathbf{F}}(m)$ for some $m \in \mathsf{max}\,\mathbf{G}(k)$. We have

$$\mathsf{max}\,\mathbf{G}(k) = \mathsf{sons}_{\mathbf{G}(k)}(0_{\mathbf{G}(k)}) \qquad \text{and} \qquad \mathsf{sons}_{\mathbf{G}(k)}(0_{\mathbf{G}(k)}) = \mathsf{sons}_{\mathbf{H}}(0_{\mathbf{G}(k)})$$

by (d1) and by (a) of Proposition 30, so it follows from (f4) and from the inductive hypothesis that $\mathsf{height}_{\mathbf{H}}(m)$ is odd and then, using (f4) and the inductive hypothesis again, we have $m \notin \big\{0_{\mathbf{G}(j)} : j \in k+1\big\}$. Let us show that $m \notin \big\{0_{\mathbf{G}(j)} : j \in |A|\big\}$. If not, then

$$m \in \big\{0_{\mathbf{G}(j)} : j \in |A| \setminus (k+1)\big\}, \qquad \text{so} \qquad m \in \bigcup\big\{M_j : j \in |A| \setminus (k+1)\big\}$$

by (f4). Then it follows from (f2) that $m \in (M_{k+1})\!\downarrow_{\mathbf{F}}$ — that is, $m \geqslant_{\mathbf{F}} t$ for some $t \in M_{k+1}$. Since $u \in \mathsf{sons}_{\mathbf{F}}(m)$, then $u >_{\mathbf{F}} m$, therefore $u >_{\mathbf{F}} t$. This contradicts (f1) because $t, u \in M_{k+1}$. Now it follows from (d4) and (a) of Proposition 30 that $\mathsf{sons}_{\mathbf{F}}(m) = \mathsf{sons}_{\mathbf{H}}(m)$, so $u \in \mathsf{sons}_{\mathbf{H}}(m)$. Then $\mathsf{height}_{\mathbf{H}}(u) \in \{2n : n \in \omega\}$ because $\mathsf{height}_{\mathbf{H}}(m)$ is odd.

Now suppose that $p \in X \setminus A$; we must find a strictly increasing function $f_p{:}\,\omega \to \omega$ that satisfies (♦). First, using (d) of Corollary 10, for each $n \in \omega$, we define $m(p, n)$ to be the node in $\mathsf{scope}_{\mathbf{H}}(p)$ such that $\mathsf{height}_{\mathbf{H}}\big(m(p, n)\big) = 2n + 1$. Using (g3) we have

$$\forall n \in \omega \ \big[m(p, n) \in \mathsf{scope}_{\mathbf{F}}(p)\big]. \tag{19}$$

Next, using (g2), we can define

$$f_p(n) \coloneqq \mathsf{height}_{\mathbf{F}}\big(m(p,n)\big) \quad \text{for every } n \in \omega;$$

then $f_p{:}\omega \to \omega$ by (a) of Corollary 10. If $n'' > n'$, then $m(p,n'') >_{\mathbf{H}} m(p,n')$ (because $\mathsf{scope}_{\mathbf{H}}(p)$ is a chain in $\mathbf{H}$ by (c) of Corollary 10), so $m(p,n'') >_{\mathbf{F}} m(p,n')$ by (g2) and (b) of Lemma 30. This implies that f_p is strictly increasing. Now, using (f4) and (g4), for every $n \in \omega$, we have

$$m(p,n) \notin \big\{0_{\mathbf{G}(i)} : i \in |A|\big\}, \quad \text{so} \quad \mathsf{sons}_{\mathbf{H}}\big(m(p,n)\big) = \mathsf{sons}_{\mathbf{F}}\big(m(p,n)\big)$$

by (d4) and (a) of Proposition 30. Then (g3) and (g1) imply

$$\mathsf{shoot}_{\mathbf{H}}\big(m(p,n)\big) \gg \mathsf{shoot}_{\mathbf{F}}\big(m(p,n)\big) \qquad \text{for all } n \in \omega. \tag{20}$$

To complete the proof it remains to verify (♦); suppose that

$$U \in \mathsf{nbhds}(p,X), \quad n \in \omega, \quad \text{and} \quad f_p(n) \in \mathsf{span}_{\mathbf{F}}(p,U).$$

The last formula means that $f_p(n) = \mathsf{height}_{\mathbf{F}}(v)$ for some $v \in \mathsf{scope}_{\mathbf{F}}(p)$ such that $\mathsf{shoot}_{\mathbf{F}}(v) \gg \{U\}$. Then $v = m(p,n)$ by (d) of Corollary 10, by (19), and by definition of $f_p(n)$. It follows that

$$\mathsf{shoot}_{\mathbf{F}}\big(m(p,n)\big) \gg \{U\}, \quad \text{so} \quad \mathsf{shoot}_{\mathbf{H}}\big(m(p,n)\big) \gg \{U\}$$

by (20). Then (g3) implies $\mathsf{shoot}_{\mathbf{H}}\big(m(p,n)\big) \gg \{U \setminus A\}$, therefore

$$2n+1 \;=\; \mathsf{height}_{\mathbf{H}}\big(m(p,n)\big) \;\in\; \mathsf{span}_{\mathbf{H}}(p, U \setminus A)$$

by definition of $\mathsf{span}_{\mathbf{H}}(p, U \setminus A)$. □

7 New examples of spaces with a π-tree

Recall that $\mathcal{N}$ is the Baire space, $\mathcal{R}_{\mathcal{S}}$ is the Sorgenfrey line, and $\mathcal{I}_{\mathcal{S}} \coloneqq \mathcal{R}_{\mathcal{S}} \setminus \mathbb{Q}$ is the irrational Sorgenfrey line.

Lemma 24.

(a) *$\mathcal{N}$ has a π-tree $\mathbf{F}$ such that* $\mathsf{cofin}\,\omega \gg \mathsf{spectrum}_{\mathbf{F}}(\mathcal{N})$.

(b) *$\mathcal{R}_{\mathcal{S}}$ has a π-tree $\mathbf{G}$ such that* $\mathsf{cofin}\,\omega \gg \mathsf{spectrum}_{\mathbf{G}}(\mathcal{R}_{\mathcal{S}})$.

(c) *If $X \subseteq \mathcal{R}_{\mathcal{S}}$ and $\mathcal{R}_{\mathcal{S}} \setminus X$ is at most countable, then X has a π-tree $\mathbf{H}$ such that* $\mathsf{cofin}\{2n+1 : n \in \omega\} \gg \mathsf{spectrum}_{\mathbf{H}}(X)$.

Proof. Part (a) follows from (b) of Lemma 6 and Example 12; part (b) can be derived from the proof of Lemma 3.6 in [2]; part (c) follows from part (b) and Corollary 21. □

Using the above lemma and Theorem 17 we obtain the following statement:

Corollary 25. *Suppose that $1 \leqslant |A| \leqslant \omega$ and for each $\alpha \in A$,*

$$\text{either} \qquad X_\alpha = \mathcal{N} \qquad \text{or} \qquad X_\alpha \subseteq \mathcal{R}_{\mathcal{S}} \;\; \text{with} \;\; |\mathcal{R}_{\mathcal{S}} \setminus X_\alpha| \leqslant \omega.$$

Then the product $\prod_{\alpha\in A} X_\alpha$ has a π-tree. □

Corollary 26.

(a) *$\mathcal{R}_{\mathcal{S}}{}^n$ and $\mathcal{I}_{\mathcal{S}}{}^n$ have a π-tree for all $n \in \omega \setminus \{0\}$.*

(b) *$\mathcal{R}_{\mathcal{S}}{}^\omega$ and $\mathcal{I}_{\mathcal{S}}{}^\omega$ have a π-tree.* □

Note that if $X \subseteq \mathcal{N}$ with $|\mathcal{N} \setminus X| \leqslant \omega$, then X is homeomorphic to $\mathcal{N}$ (this can be easily derived from the Alexandrov-Urysohn characterization of the Baire space and from the characterization of its Polish subspaces — see Theorems 3.11 and 7.7 in [4]). Notice also that $\mathcal{N}^n$ is homeomorphic to $\mathcal{N}$ for all $n \in \omega \setminus \{0\}$ and $\mathcal{N}^\omega$ is also homeomorphic to $\mathcal{N}$.

Corollary 27. *If a space X has a π-tree, then $X \times \mathcal{N}$, $X \times \mathcal{R}_{\mathcal{S}}$, and $X \times \mathcal{R}_{\mathcal{S}}{}^\omega$ also have a π-tree.*

Proof. This statement follows from Corollary 15 and Lemma 24. □

8 Appendix. The foliage hybrid operation

In the proofs of Lemma 19 and Proposition 20 we employ the foliage hybrid operation, which was introduced in [1]. For completeness of exposition we list here definitions and results that we use. The definition of graft, which we give below, slightly differs from the definition of graft in [1], but these two definitions are easily seen to be equivalent. The same can be said about our definition of $\mathsf{hybrid}(\mathcal{T},\gamma)$, see details in [1, Remark 20]. To ease comprehension of notions from Definition 29, you can look at pictures that illustrate this definition in [10].

Notation 28.

- $\forall x \neq y \in A\ \varphi(x,y) \quad :\longleftrightarrow \quad \forall x,y \in A\ \big[x \neq y \to \varphi(x,y)\big]$;
- $x \parallel_{\mathcal{P}} y \quad :\longleftrightarrow \quad x \not\leqslant_{\mathcal{P}} y$ and $x \not\geqslant_{\mathcal{P}} y$.

Definition 29 (Definitions 15, 17, 19, 25–27 and Remark 20 in [1])**.** Suppose that $\mathcal{T},\mathcal{G}$ are trees and $\mathbf{F},\mathbf{G}$ are nonincreasing foliage trees.

- $\mathcal{G}$ is a **graft** for $\mathcal{T} \quad :\longleftrightarrow$
  - $|\mathsf{nodes}\,\mathcal{G}| > 1$,
  - $\mathcal{G}$ has the least node,
  - $\mathsf{nodes}\,\mathcal{G} \cap \mathsf{nodes}\,\mathcal{T} = \{0_{\mathcal{G}}\} \cup \mathsf{max}\,\mathcal{G}$, and
  - $\forall x,y \in \mathsf{nodes}\,\mathcal{G} \cap \mathsf{nodes}\,\mathcal{T}\ [x <_{\mathcal{G}} y \leftrightarrow x <_{\mathcal{T}} y]$.
- If $\mathcal{G}$ is a graft for $\mathcal{T}$, then:
  - $\mathsf{implant}\,\mathcal{G} := \mathsf{nodes}\,\mathcal{G} \setminus \big(\{0_{\mathcal{G}}\} \cup \mathsf{max}\,\mathcal{G}\big)$;
  - $\mathsf{explant}(\mathcal{T},\mathcal{G}) := (0_{\mathcal{G}})\!\downarrow_{\mathcal{T}} \setminus (\mathsf{max}\,\mathcal{G})\!\downarrow_{\mathcal{T}}$.
- γ is a **consistent** family of grafts for $\mathcal{T} \quad :\longleftrightarrow$
  - $\forall \mathcal{G} \in \gamma\ [\,\mathcal{G}$ is a graft for $\mathcal{T}\,]$,
  - $\forall \mathcal{D} \neq \mathcal{E} \in \gamma\ [\,\mathsf{implant}\,\mathcal{D} \cap \mathsf{implant}\,\mathcal{E} = \varnothing\,]$, and
  - $\forall \mathcal{D} \neq \mathcal{E} \in \gamma\ \big[\,0_{\mathcal{D}} \parallel_{\mathcal{T}} 0_{\mathcal{E}}$ or $0_{\mathcal{D}} \in (\mathsf{max}\,\mathcal{E})\!\downarrow_{\mathcal{T}}$ or $0_{\mathcal{E}} \in (\mathsf{max}\,\mathcal{D})\!\downarrow_{\mathcal{T}}\,\big]$.
- If γ is a consistent family of grafts for $\mathcal{T}$, then:
  - $\mathsf{support}(\mathcal{T},\gamma) := \mathsf{nodes}\,\mathcal{T} \setminus \bigcup\limits_{\mathcal{G}\in\gamma} \mathsf{explant}(\mathcal{T},\mathcal{G})$;
  - $\mathsf{hybrid}(\mathcal{T},\gamma) :=$ the pair $(H,<)$ (actually, a tree) such that
    - ✓ $H := \mathsf{support}(\mathcal{T},\gamma) \cup \bigcup\limits_{\mathcal{G}\in\gamma} \mathsf{implant}\,\mathcal{G}$ and
    - ✓ $<\ :=$ the transitive closure of relation $(<_{\mathcal{T}} \cup \bigcup\limits_{\mathcal{G}\in\gamma} <_{\mathcal{G}}) \cap (H \times H)$.
- $\mathbf{G}$ is a **foliage graft** for $\mathbf{F} \quad :\longleftrightarrow$
  - $\mathbf{G}$ is nonincreasing,
  - $\mathsf{skeleton}\,\mathbf{G}$ is a graft for $\mathsf{skeleton}\,\mathbf{F}$,
  - $\mathbf{G}_{0_{\mathbf{G}}} \subseteq \mathbf{F}_{0_{\mathbf{G}}}$, and
  - $\forall m \in \mathsf{max}\,\mathbf{G}\ [\mathbf{G}_m = \mathbf{F}_m]$.
- If $\mathbf{G}$ is a foliage graft for $\mathbf{F}$, then
  - $\mathsf{cut}(\mathbf{F},\mathbf{G}) := \mathbf{F}_{0_{\mathbf{G}}} \setminus \mathbf{G}_{0_{\mathbf{G}}}$.
- φ is a **consistent** family of foliage grafts for $\mathbf{F} \quad :\longleftrightarrow$
  - $\forall \mathbf{G} \in \varphi\ [\mathbf{G}$ is a foliage graft for $\mathbf{F}]$,
  - $\forall \mathbf{D} \neq \mathbf{E} \in \varphi\ [\mathsf{skeleton}\,\mathbf{D} \neq \mathsf{skeleton}\,\mathbf{E}]$, and
  - $\{\mathsf{skeleton}\,\mathbf{G} : \mathbf{G} \in \varphi\}$ is a consistent family of grafts for $\mathsf{skeleton}\,\mathbf{F}$.
- If φ is a consistent family of foliage grafts for $\mathbf{F}$, then:

➢ $\mathsf{loss}(\mathbf{F},\varphi) := \bigcup_{\mathbf{G}\in\varphi} \mathsf{cut}(\mathbf{F},\mathbf{G})$;

➢ $\mathsf{fol.hybr}(\mathbf{F},\varphi)$:= the **foliage hybrid** of $\mathbf{F}$ and φ := the foliage tree $\mathbf{H}$ such that

✓ $\mathsf{skeleton}\,\mathbf{H} := \mathsf{hybrid}\big(\mathsf{skeleton}\,\mathbf{F}, \{\mathsf{skeleton}\,\mathbf{G} : \mathbf{G}\in\varphi\}\big)$ and

✓ $\mathbf{H}_x := \begin{cases} \mathbf{G}_x \smallsetminus \mathsf{loss}(\mathbf{F},\varphi), & \text{if } x\in\mathsf{implant}\,\mathbf{G} \text{ for some } \mathbf{G}\in\varphi; \\ \mathbf{F}_x \smallsetminus \mathsf{loss}(\mathbf{F},\varphi), & \text{otherwise.} \end{cases}$

Lemma 30 (Lemma 21 and Proposition 23 in [1]). *Suppose that γ is a consistent family of grafts for a tree $\mathcal{T}$, $\mathcal{H} = \mathsf{hybrid}(\mathcal{T},\gamma)$, and $\mathcal{G}\in\gamma$.*

(a) $\mathsf{nodes}\,\mathcal{G}\subseteq\mathsf{nodes}\,\mathcal{H}$ *and* $\forall x,y\in\mathsf{nodes}\,\mathcal{G}\ [x<_{\mathcal{H}} y \leftrightarrow x<_{\mathcal{G}} y]$.

(b) $\mathsf{support}(\mathcal{T},\gamma) = \mathsf{nodes}\,\mathcal{H}\cap\mathsf{nodes}\,\mathcal{T}$ *and* $\forall x,y\in\mathsf{support}(\mathcal{T},\gamma)\ [x<_{\mathcal{H}} y \leftrightarrow x<_{\mathcal{T}} y]$.

(c) *For each* $x\in\mathsf{nodes}\,\mathcal{H}$,

$$\mathsf{sons}_{\mathcal{H}}(x) = \begin{cases} \mathsf{sons}_{\mathcal{G}}(x), & \text{if } x\in\{0_{\mathcal{G}}\}\cup\mathsf{implant}\,\mathcal{G} \text{ for some } \mathcal{G}\in\gamma; \\ \mathsf{sons}_{\mathcal{T}}(x), & \text{otherwise } \big(\text{i.e., when } x\in\mathsf{support}(\mathcal{T},\gamma)\smallsetminus\{0_{\mathcal{G}}:\mathcal{G}\in\gamma\}\big). \end{cases}$$

Lemma 31 (Lemma 30 in [1]). *Suppose that $\mathbf{F}$ is a Baire foliage tree on a space X and φ is a consistent family of foliage grafts for $\mathbf{F}$ such that every $\mathbf{G}$ in φ is ω-branching, locally strict, open in X, has bounded chains, and has $\mathsf{height}\,\mathbf{G}\leqslant\omega$. Then the foliage hybrid of $\mathbf{F}$ and φ is a Baire foliage tree on $X\smallsetminus\mathsf{loss}(\mathbf{F},\varphi)$.*

Definition 32 (Definitions 31, 33 in [1]). Suppose that $\mathbf{H},\mathbf{F}$ are nonincreasing foliage trees and $\mathbf{G}$ is a foliage graft for $\mathbf{F}$.

✎ **H shoots into F** $:\longleftrightarrow$ $\forall p\in\mathsf{flesh}\,\mathbf{H}\ \forall y\in\mathsf{scope}_{\mathbf{F}}(p)\ \exists x\in\mathsf{scope}_{\mathbf{H}}(p)\ \big[\mathsf{shoot}_{\mathbf{H}}(x)\gg\mathsf{shoot}_{\mathbf{F}}(y)\big]$.

✎ **G preserves shoots** of **F** $:\longleftrightarrow$

for each $p\in\mathsf{flesh}\,\mathbf{G}$ and for each $y\in\mathsf{scope}_{\mathbf{F}}(p)\cap\big(\{0_{\mathbf{G}}\}\cup\mathsf{explant}(\mathbf{F},\mathbf{G})\big)$

there is $x\in\mathsf{scope}_{\mathbf{G}}(p)\cap\big(\{0_{\mathbf{G}}\}\cup\mathsf{implant}\,\mathbf{G}\big)$ such that $\big[\mathsf{shoot}_{\mathbf{G}}(x)\gg\mathsf{shoot}_{\mathbf{F}}(y)\big]$.

Lemma 33 (Lemma 34 in [1]). *Suppose that $\mathbf{F}$ is a nonincreasing foliage tree, φ is a consistent family of foliage grafts for $\mathbf{F}$, the foliage hybrid of $\mathbf{F}$ and φ has nonempty leaves, and each $\mathbf{G}\in\varphi$ preserves shoots of $\mathbf{F}$. Then the foliage hybrid of $\mathbf{F}$ and φ shoots into $\mathbf{F}$.*

Lemma 34 (Lemma 32 in [1]). *Suppose that a foliage tree $\mathbf{H}$ shoots into a foliage tree $\mathbf{F}$ and $\mathbf{F}$ grows into a space X. Then $\mathbf{H}$ grows into the subspace $X\cap\mathsf{flesh}\,\mathbf{H}$ of X.*

References

[1] M. Patrakeev, The complement of a σ-compact subset of a space with a π-tree also has a π-tree, *Topology and its Applications*, the special issue dedicated to the 120th anniversary of P.S. Alexandroff, accepted for publication in 2016. Preprint: `https://arxiv.org/abs/1512.02458` .

[2] M. Patrakeev, Metrizable images of the Sorgenfrey line, *Topology Proceedings* vol. 45 (2015) 253–269.

[3] A. V. Arhangel'skii, *Open and close-to-open mappings. Relations among spaces*, Trudy Moskov Mat. Obšč. 1966, no. 15, 181–223.

[4] A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer, 1994.

[5] I. Juhász, *Cardinal functions in topology: ten years later,* MC tracts, vol. 123, 1980.

[6] E. K. van Douwen, W. F. Pfeffer, Some properties of the Sorgenfrey line and related spaces, *Pacific J. Math.*, vol. 81, no. 2 (1979), 371–377.

[7] T. Jech, *Set Theory*, 2nd Edition, Springer, 1996.

[8] K. Kunen, *Set Theory*, North-Holland, 1980.

[9] K. P. Hart, J. Nagata, and J. E. Vaughan, eds., *Encyclopedia of general topology*, Elsevier, Amsterdam, 2004.

[10] M. Patrakeev, *The foliage hybrid operation*, slides from the talk at Alexandroff Readings, May 22–26, 2016, Moscow, Russia; available at `https://arxiv.org/src/1512.02458v5/anc` .